



FIELD RESEARCH BOOK

2018 Edition - Huxley Learning Center



Throughout the state in any given year, our products are put to the test in various research trials. These trials allow us to gain insights to help complete a year-long story of product performance, agronomic characteristics, and weather patterns.

We spend months planning out our research trials, but as you know, they are vulnerable to extreme weather patterns and unforeseen challenges. Across the state, we saw extreme drought in the south, heavy rain across the north, a new soybean insect in the west, and a corn disease creeping in from the east. With conditions like this, we often see research results that may not match data from prior years, but we can use these challenges as an opportunity to evaluate, learn, and provide better insights for the future.

For example, due to much of the available nitrogen being moved away from the ideal zone, we didn't see the yield response to planting populations that we typically see in high yield environments. On the flip side, we were able to evaluate how products handle nitrogen deficiency. This is just one scenario provided throughout our agronomic reports. I hope you enjoy all the information and find it insightful.

I'd also like to recognize my Bayer Crop Science colleagues who contributed to the research in this booklet:

Charles Boateng - Huxley

Jarod Jackson, Eastern Iowa

Matt Nelson - Atlantic

Brett Schafer, Western Iowa

Craig Lamoureux - Storm Lake

John Cantwell, Iowa

Chuck Kolbet - Marble Rock

Doug Doty - Victor

We appreciate your relationship in 2018 and look forward to working with you in 2019!

Brent Schwenneker

Huxley Learning Center



Table of Contents

5	Management Practices for Optimizing Yield and Productivity in Soybean
7	Soybean Yield Response to Seeding Rate
11	Management of Early-season Stressors in Soybeans
15	Yield Observations When Shifting To Earlier Maturity Group Soybeans
17	Management Practices for Optimizing Yield and Productivity in Corn
21	Yield Response of Corn Products to Seeding Rate
25	Characterization of Corn Products for Their Response to Nitrogen Fertilization
29	Effects of Nitrogen Rate on Corn Yield Potential
32	Effect of Fungicide on Yield and Plant Health
35	2018 FFA Fantasy Farming Challenge in Iowa
38	Comparison of Row Spacing by Management Practice
41	Comparison of Corn Row Spacing and Seeding Rate - Storm Lake, IA
44	The Effects Of Row Spacing And Seeding Rates On Corn Yield Potential
46	Tillage Systems in Corn and Soybean Production
49	Cover Cropping and Tillage Systems in Soybean Production
51	Using 2018 Corn Rootworm Beetle Counts To Help Evaluate the Risk of an Infestation for 2019

How to Use This Book

The reports in this book are arranged by section: soybean, corn and agronomics. Each report is also tagged with one of these icons to quickly show you what it's about.





Management Practices for Optimizing Yield and Productivity in Soybean

Trial Objective

- Obtaining higher yields in soybean production involves the efficient and sustainable use of farm resources and management practices.
- There are several inputs and practices that growers use each year that ultimately impact yield and profitability. Therefore, the objective of this study was to evaluate the economic impact of these inputs and practices.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Huxley, IA	Clay loam	Corn	No tillage	5/10/2018	10/17/2018	60	140K

- A 2.0 maturity group soybean product was planted in 200-ft long strips.
- The trial was carried out in 30-inch row spacing, six rows per treatment, with two replications.
- Overall disease incidence and severity were low for this trial/location.
- The management practices tested were two seed treatments, a nitrogen side-dress application, and a fungicide application. These practices were compared in incremental stair-step treatments (Table 1).
- Acceleron® Seed Applied Solutions STANDARD is a fungicide and insecticide seed treatment.
- ILeVO® is a systemic soybean seed treatment for protection against early-season damage caused by pathogenic nematodes and *Fusarium virguliforme*, which causes Sudden Death Syndrome (SDS).
- Delaro® 325 SC fungicide was applied at R3 for the fungicide treatment.
- 32% urea and ammonium nitrate (UAN) was applied at the R3 growth stage to deliver 40 lb/acre of nitrogen.

Table 1. Treatments used in the trial with associated costs. Treatment cost does not include the price of seed.

Treatments	Input	Treatment Cost (\$/acre)
UTC	Untreated	\$ -
A	Acceleron® Seed Applied Solutions STANDARD (A)	\$7.00
A+I	ILeVO® seed treatment (I)	\$19.00
AI+F	Foliar fungicide application at R3 (F)	\$41.00
AIF+N	Side dress 32% UAN at R3 (N)	\$50.20

Understanding the Results

- All inputs improved yields over the untreated plot (UTC).
- The addition of ILeVO® seed treatment provided the largest yield response of all of the treatments.
- The addition of a fungicide application had a minimal effect on yield over the previous treatment and the addition of nitrogen did not provide a yield response over the previous treatment.



Management Practices for Optimizing Yield and Productivity in Soybean




Figure 1. Productivity of soybean under the different management systems. Soybean grain price was set at \$7.50/bu.

- The yield gained by some of the systems (treatments A and AIFN) were not high enough to provide an economic incentive over the untreated plot (UTC).
- A system with just the two seed treatments provided the most economical input in this trial.

What Does This Mean For Your Farm?

- During the 2018 growing season, the research site experienced wet and rainy conditions in May and June followed by a dry July. Such conditions can result in poor plant health and may explain why the Acceleron® Seed Applied Solutions STANDARD with ILeVO® (AI) treatment was the most profitable.
- Growers should consider performing small-scale trials on their fields to understand how their management systems impact their operations economically.

Legal Statements

The information discussed in this report is from a single-site, replicated demonstration trial. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

FOR SOYBEANS, EACH ACCELERON® SEED APPLIED SOLUTIONS OFFERING is a combination of separate individually registered products containing the active ingredients: **BASIC** Offering: metalaxyl, flupyrroxad, and pyraclostrobin. **STANDARD** Offering: metalaxyl, flupyrroxad, pyraclostrobin, and imidacloprid. **STANDARD FN** Offering: metalaxyl, flupyrroxad, pyraclostrobin, and tiozazafen. **ELITE** Offering: metalaxyl, flupyrroxad, pyraclostrobin, imidacloprid, and tiozazafen. **Upstream Treatment Offerings Only:** Acceleron® B-200 SAT is included seamlessly in the Acceleron® Seed Applied Solutions **STANDARD FN** and **ELITE** tiers.

The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Not all products are approved in all states. Please visit <http://acceleronsas.com/stateapprovals> regarding the approval status of products containing NemaStrike™ Technology for application to seeds in your state.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. ILeVO® is a trademark of BASF Corporation. Acceleron®, Delaro® and NemaStrike™ are trademarks of Bayer Group. Asgrow and the A Design®, Asgrow®, DEKALB and Design® and DEKALB® are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. ©2018 Bayer Group. All rights reserved.181129143915 112918TAM





Soybean Yield Response to Seeding Rate

Trial Objective

- A critical agronomic decision is the choice of soybean products and their associated optimum seeding rate. Though soybeans compensate well at different plant populations, soybean products respond differently and the search for the optimum seeding rate continues.
- The objective of this study was to determine the yield potential of Asgrow® brand soybean products at different seeding rates.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seed/acre)
Huxley, IA	Clay loam	Corn	No tillage	05/09/2016	10/17/2016	60	100K, 140K, 180K

- Fifteen Asgrow® brand soybean products ranging from 1.8 to 3.6 maturity group (MG) were planted at 100,000, 140,000, and 180,000 seeds/acre.
- The trial was planted in 30-inch spacing, 4 rows per product per seeding rate, and 200-ft long plots.
- Weed management consisted of a rye burndown with Roundup PowerMAX® herbicide and a post-emergence application of XtendiMax® herbicide with VaporGrip® Technology, Warrant® herbicide, and Roundup PowerMAX®.

Understanding the Results

- The decrease in plant population across the trial between the V4 and R8 growth stages was partly due to wet and rainy conditions at the research site in May and June, followed by a dry July. Such conditions result in poor seedling health and reduced crop stands.
- Seeding rate did not have an impact on lodging in most of the products tested, especially in the late MG products (Table 1 and 2). In the products in which there was an association (particularly in the early MG), lodging increased with increasing seeding rate.
- In both the early and late MGs, yield increased as seeding rate increased in most of the products. In the other products, the 140,000 seeds/acre seeding rate produced the highest yields (Figure 1 and 2).
- In both MGs, the biggest yield increase was from the 100,000 to 140,000 seeds/acre rate increase (Figure 1 and 2).



Soybean Yield Response to Seeding Rate

Table 1. Performance of early MG Asgrow® brand soybean products at different seeding rates. Early stand count was taken at the V4 growth stage. Harvest population and lodging score were taken after the R8 growth stage. Lodging score was based on a 1-9 scale with 1 = 100% erect and 9 = 100% flat.

Asgrow® Brand	Seeding Rate (1000 seeds/acre)	Early Stand Count (1000 seeds/acre)	Harvest Population (1000 seeds/acre)	Lodging Score	Grain Moisture (%)	Average Yield (bu/acre)	Average Yield Ranking
AG18X9	100	81.5	84.5	1	11.1	62.2	7
	140	105	101.5	1	11.5		
	180	149	123.5	1	11.2		
AG20X9	100	90.5	85.5	1.5	12.7	68.0	3
	140	123.5	108	3	13.0		
	180	149.5	129	5	13.3		
AG21X9	100	88	78	2	12.6	70.1	2
	140	124.5	107	3.5	13.0		
	180	155	132	5.5	13.0		
AG22X9	100	83	77.5	2	12.2	70.6	1
	140	113.5	107	3	12.5		
	180	142.5	124	5.5	12.6		
AG23X9	100	86.5	86	1.5	11.3	63.8	6
	140	120.5	114	2.5	11.5		
	180	145	132.5	5	11.5		
AG24X9	100	75.5	82	1	11.7	64.2	5
	140	109.5	99	1	11.7		
	180	133	119.5	2	11.6		
AG25X9	100	68.5	73.5	1	11.4	66.7	4
	140	100.5	95.5	2	11.5		
	180	128.5	115.5	2	11.6		




Figure 1. Average yields of early MG Asgrow® brand soybean products at three different seeding rates.



Soybean Yield Response to Seeding Rate

Table 2. Performance of late MG Asgrow® brand soybean products at different seeding rates. The early stand count was taken at the V4 growth stage. Harvest population and lodging score were taken after the R8 growth stage. Lodging score was based on a 1-9 scale with 1 = 100% erect and 9 = 100% flat.

Asgrow® Brand	Seeding Rate (1000 seeds/acre)	Early Stand Count (1000 seeds/acre)	Harvest Population (1000 seeds/acre)	Lodging Score	Grain Moisture (%)	Average Yield (bu/acre)	Average Yield Rating
AG26X8	100	84.5	82	1	11.4	62.3	5
	140	115.5	102.5	1	11.3		
	180	143.5	124	1	11.5		
AG27X9	100	88	83.5	1	11.5	57.2	6
	140	121.5	122	1.5	11.4		
	180	152.5	126.5	2.5	11.4		
AG28X9	100	81	81	1	12.0	66.1	3
	140	116	108.5	1	11.8		
	180	139.5	121	1	11.8		
AG29X9	100	86.5	88	1	12.6	74.1	1
	140	115	112.5	1	12.5		
	180	155	142.5	1.5	12.5		
AG30X9	100	90	87	1	12.9	64.6	4
	140	123.5	112	1	13.0		
	180	149.5	141.5	1	12.9		
AG31X9	100	73.5	77	1	13.5	56.8	7
	140	92.5	95	1	12.4		
	180	115.5	106	1	13.1		
AG33X8	100	83	80.5	1	13.9	56.3	8
	140	115.5	112.5	1	13.9		
	180	156	143.5	1	14.1		
AG36X6	100	80	83	1	12.9	71.7	2
	140	115	104	1	12.8		
	180	153	134.5	2	12.5		




Figure 2. Average yields of late MG Asgrow® brand soybean products at three different seeding rates.



Soybean Yield Response to Seeding Rate

What Does This Mean for Your Farm?

- With the current soybean seed cost at \$62/bag and assuming the soybean grain was sold at \$7.50/bu, a minimum of a 2.4 bu/acre yield increase is needed to justify a 40,000 seeds/acre seeding rate increase.
- Thus, for all products in the early MG (except for AG20X9 and AG21X9), 180,000 seeds/acre was the highest yielding and most profitable seeding rate (Figure 1).
- For the late MG, the median seeding rate of 140,000 seeds/acre was the most profitable, except for AG30X9, in which the 180,000 seeds/acre rate was both the highest yielding and most profitable (Figure 2).
- This is our fourth year of research into germplasm response to seeding rate. Results from the 2015-2017 trials were product-specific and did not provide a consistent trend in response to seeding rate.¹ The 2018 results are consistent across almost all products tested, providing a good benchmark for future decision making.
- Growers should consult their trusted agronomists and dealers in choosing the best products for their operation.

Source

¹Monsanto Learning Center at Huxley, Iowa Demonstration Reports, 2015-017.

Legal Statements

XtendiMax® herbicide with VaporGrip® Technology is part of the Roundup Ready® Xtend Crop System and is a restricted use pesticide. ALWAYS READ AND FOLLOW DIRECTIONS FOR USE ON PESTICIDE LABELING. It is a violation of federal and state law to use any pesticide product other than in accordance with its labeling. XtendiMax® herbicide with VaporGrip® Technology and products with XtendFlex® Technology may not be approved in all states and may be subject to use restrictions in some states. Check with your local product dealer or representative or U.S. EPA and your state pesticide regulatory agency for the product registration status and additional restrictions in your state. For approved tank-mix products and nozzles visit [XtendiMaxApplicationRequirements.com](#)

Certain statements contained in this communication are “forward-looking statements,” such as statements concerning the company’s anticipated financial results, current and future product performance, regulatory approvals, business and financial plans and other non-historical facts. These statements are based on current expectations and currently available information. However, since these statements are based on factors that involve risks and uncertainties, the company’s actual performance and results may differ materially from those described or implied by such forward-looking statements. Factors that could cause or contribute to such differences include, among others: the risk of disruptions or uncertainties related to the transaction with Bayer Aktiengesellschaft and the risk that the expected benefits of the transaction will not be realized in the anticipated time frame or at all; continued competition in seeds, traits and agricultural chemicals; the company’s exposure to various contingencies, including those related to intellectual property protection, regulatory compliance and the speed with which approvals are received, and public understanding and acceptance of our biotechnology and other agricultural products; the success of the company’s research and development activities; the outcomes of major lawsuits; developments related to foreign currencies and economies; fluctuations in commodity prices; compliance with regulations affecting our manufacturing; the accuracy of the company’s estimates related to distribution inventory levels; the levels of indebtedness, continued availability of capital and financing and rating agency actions; the company’s ability to fund its short-term financing needs and to obtain payment for the products that it sells; and the effect of weather conditions, natural disasters, accidents, and security breaches, including cyber security incidents, on the agriculture business or the company’s facilities. Undue reliance should not be placed on these forward-looking statements, which are current only as of the date of this communication. The company disclaims any current intention or obligation to update any forward-looking statements or any of the factors that may affect actual results.

This information is for educational purposes only and is not an offer to sell Roundup Xtend™ with VaporGrip™ Technology. These products are not yet registered or approved for sale or use anywhere in the United States.

Commercialization is dependent on multiple factors, including successful conclusion of the regulatory process. **The information presented herein is provided for educational purposes only, and is not and shall not be construed as an offer to sell, or a recommendation to use, any unregistered pesticide for any purpose whatsoever.** It is a violation of federal law to promote or offer to sell an unregistered pesticide.

Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto’s Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship.

ALWAYS READ AND FOLLOW DIRECTIONS FOR USE ON PESTICIDE LABELING. IT IS A VIOLATION OF FEDERAL AND STATE LAW to use any pesticide product other than in accordance with its labeling. NOT ALL formulations of dicamba or glyphosate are approved for in-crop use with Roundup Ready 2 Xtend® soybeans. ONLY USE FORMULATIONS THAT ARE SPECIFICALLY LABELED FOR SUCH USES AND APPROVED FOR SUCH USE IN THE STATE OF APPLICATION. Contact the U.S. EPA and your state pesticide regulatory agency with any questions about the approval status of dicamba herbicide products for in-crop use with Roundup Ready 2 Xtend® soybeans.

Roundup Ready 2 Xtend® soybeans contains genes that confer tolerance to glyphosate and dicamba. Glyphosate will kill crops that are not tolerant to glyphosate. Dicamba will kill crops that are not tolerant to dicamba. Contact your seed brand dealer or refer to the Technology Use Guide for recommended weed control programs.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Warrant® Herbicide is not registered in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Roundup Xtend™ is a trademark of Monsanto Technology LLC. Asgrow and the A Design®, Asgrow®, DEKALB and Design®, Roundup Ready 2 Xtend®, Roundup Ready®, VaporGrip®, Warrant® and XtendiMax® are registered trademarks of Bayer Group. ©2018 Bayer Group. All rights reserved. 181108081856 110918TAM





Management of Early-season Stressors in Soybean Production

Trial Objective

- To optimize yield potential, soybean growers often plant as early as possible within the recommended planting window. Such early planting predisposes seeds and young seedlings to early-season stressors, such as insects, diseases, and cool, wet soils, which can significantly affect stand establishment and overall plant and field health.
- To help alleviate some of the early planting challenges, the use of seed treatments has become an important tool in fields prone to early-season stressors. Insecticides, fungicides, and nematicides are the common components of most seed treatments, and the choice depends on the anticipated pest to be controlled.
- The objective of this study was to determine soybean product performance as influenced by two seed treatment options.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Huxley, IA	Clay loam	Corn	No tillage	05/09/2018	10/17/2018	60	140K

- Twenty-seven soybean products ranging from 1.8 to 3.7 maturity group (MG) were planted at an average of 140,000 seeds/acre.
- Each product received two types of seed treatments:
 - Base seed treatment (fungicide and insecticide)
 - Base + ILeVO® seed treatment
 - ILeVO® is a systemic soybean seed treatment for protection against early-season damage caused by pathogenic nematodes and *Fusarium virguliforme*, which causes Sudden Death Syndrome (SDS).
- The trial was planted in 30-inch row spacing, two rows per product per treatment, 200-ft long plots, and three replications.
- There was a low level of SDS incidence at the research site.
- No symptoms of soybean cyst nematode (SCN) were observed at the research site. Soil samples in 2017 showed low SCN levels.
- Weed management consisted of a rye burndown with Roundup PowerMAX® herbicide and a post-emergence application of XtendiMax® herbicide with VaporGrip® Technology, Warrant® Herbicide, and Roundup PowerMAX® herbicide.



Management of Early-season Stressors in Soybean Production

Understanding the Results

- In general, ILeVO® seed treatment did not substantially affect grain moisture content and stem lodging, but did have a remarkable affect on plant density in some products (Table 1).
- Of the 27 products tested, eight (approx. 30%) did not have a positive yield response to ILeVO® seed treatment. Of the remaining 70% that responded positively, the yield advantage ranged between 0.5 to 10.8 bu/acre (Figure 1).
- The average yield response to ILeVO® seed treatment for all products was 2.53 bu/acre; however, when looking only at the products that showed a positive yield response with ILeVO® seed treatment, the average yield response was 4.8 bu/acre (Figure 1).
- In general, products with an SDS score of 3 had the lowest yield response to ILeVO® seed treatment (Figure 2).

Table 1. Agronomic response of soybean products to seed treatments. The data represent the difference between the Base + ILeVO® treatment minus the Base treatment. Early stand count was taken at the V4 growth stage. Harvest population and lodging score were taken after the R8 growth stage. Lodging score was based on a 1-9 scale with 1 = 100% erect and 9 = 100% flat. SDS scores are the official scores from product seed guides and are based on a 1-9 scale with 1 = no disease and 9 = ≥ 80% incidence with severe symptoms.

Maturity Group	Early Stand Count (1000 seeds/acre)	Harvest Population (1000 seeds/acre)	Lodging Score	SDS Score	Grain Moisture (%)
1.8	1.0	3.7	0.0	4.0	0.2
1.8	7.7	20.3	0.7	4.0	-0.4
1.8	5.0	-4.7	0.3	4.0	0.5
2.0	1.0	-0.3	-1.0	5.0	0.4
2.0	4.7	16.7	0.3	4.0	-0.3
2.1	4.3	3.3	0.0	4.0	0.4
2.2	3.3	-1.0	-0.3	4.0	0.5
2.2	4.0	13.3	0.0	5.0	0.0
2.2	3.0	-7.0	0.0	5.0	-0.1
2.3	14.0	4.3	2.0	4.0	0.5
2.4	-5.3	-1.0	0.0	3.0	0.1
2.4	5.0	15.3	0.3	4.0	0.0
2.6	7.0	6.3	0.0	5.0	-0.3
2.6	9.0	1.3	-1.3	3.0	0.0
2.7	-0.7	4.0	0.7	5.0	0.1
2.8	-3.0	1.0	0.0	4.0	0.1
2.8	-14.3	-13.0	0.0	3.0	-0.1
2.9	2.3	-4.0	-0.3	4.0	-0.1
2.9	5.3	-5.0	0.0	4.0	0.0
3.0	3.3	0.3	0.0	5.0	-0.1
3.1	1.0	-6.0	0.0	5.0	0.4
3.1	-14.0	-5.0	0.0	4.0	0.0
3.3	1.7	-2.3	0.0	3.0	-0.3
3.3	6.7	5.0	0.0	5.0	0.1
3.3	-4.0	-7.0	-0.3	3.0	-0.2
3.5	-1.7	-0.3	0.0	6.0	0.2
3.7	3.0	-0.7	0.0	5.0	-0.3
AVG	1.8	1.4	0.0	-	0.0



Management of Early-season Stressors in Soybean Production




Figure 1. Yield advantage of ILeVO® seed treatment (base seed treatment of fungicide and insecticide components + ILeVO® seed treatment over a base seed treatment alone). The red line indicates a 1.6 bu/acre economic break-even yield for ILeVO® seed treatment.




Figure 2. Impact of a soybean product's SDS score on the yield response to ILeVO® seed treatment. SDS scores are the official scores from product seed guides and are based on a 1-9 scale with 1 = no disease and 9 = ≥ 80% incidence with severe symptoms.



Management of Early-season Stressors in Soybean Production

What Does This Mean for Your Farm?

- Several seed treatment options are available to growers and a decision should be based on the challenges of the operation and the expected economic value (ROI).
- Fungicides and insecticides should be the platform upon which other seed treatment options are based, where needed.
- Not all SDS infections produce foliar symptoms. Thus, the disease could be robbing yields without growers knowing. ILeVO® seed treatment is mainly recommended for SDS control and could be an important addition to the seed treatment platform of the operation.
- With the current soybean grain price at \$7.50/bu, a minimum of a 1.6 bu/acre yield increase was needed in this trial to offset the cost of ILeVO® seed treatment (approx. \$12/acre) (Figure 1).
- In this trial, product tolerance (an SDS score of 3) provided good control of the disease, thus ILeVO® was not warranted. Where product tolerance is lacking, ILeVO® could provide more than a 2 bu/acre economic gain for the operation (Figure 2).
- As always, growers are encouraged to conduct small-scale trials on their fields to evaluate the value of new practices to their operations. They should also consult their trusted agronomists and dealers when choosing the best seed products for their operations.

Legal Statements

The information discussed in this report is from a single-site, replicated demonstration trial. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

XtendiMax® herbicide with VaporGrip® Technology is part of the Roundup Ready® Xtend Crop System and is a restricted use pesticide. ALWAYS READ AND FOLLOW DIRECTIONS FOR USE ON PESTICIDE LABELING. It is a violation of federal and state law to use any pesticide product other than in accordance with its labeling. XtendiMax® herbicide with VaporGrip® Technology and products with XtendFlex® Technology may not be approved in all states and may be subject to use restrictions in some states. Check with your local product dealer or representative or U.S. EPA and your state pesticide regulatory agency for the product registration status and additional restrictions in your state. For approved tank-mix products and nozzles visit XtendiMaxApplicationRequirements.com

FOR SOYBEANS, EACH ACCELERON® SEED APPLIED SOLUTIONS OFFERING is a combination of separate individually registered products containing the active ingredients: BASIC Offering: metalaxyl, fluxapyroxad, and pyraclostrobin. STANDARD Offering: metalaxyl, fluxapyroxad, pyraclostrobin, and imidacloprid. STANDARD FN Offering: metalaxyl, fluxapyroxad, pyraclostrobin, and tioxazafen. ELITE Offering: metalaxyl, fluxapyroxad, pyraclostrobin, imidacloprid, and tioxazafen. Upstream Treatment Offerings Only: Acceleron® B-200 SAT is included seamlessly in the Acceleron® Seed Applied Solutions STANDARD FN and ELITE tiers.

The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Not all products are approved in all states. Please visit <http://acceleronsas.com/stateapprovals> regarding the approval status of products containing NemaStrike™ Technology for application to seeds in your state.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Warrant® Herbicide is not registered in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. ILeVO® is a trademark of BASF Corporation. Acceleron®, Asgrow and the A Design®, Asgrow®, DEKALB and Design® and DEKALB®, NemaStrike™, Roundup PowerMAX®, VaporGrip®, Warrant® and XtendiMax® are trademarks of Bayer Group. ©2019 Bayer Group. All rights reserved. 181129144802 112918TAM





Yield Observations When Shifting to Earlier Maturity Group Soybeans

Trial Objective

- We continue to see a trend of growers planting earlier maturity group (MG) soybeans in Iowa and managing them at a higher level with seed treatments and foliar applications of fungicide and insecticide.
- This trend, dubbed “MG shift”, is becoming increasingly important in some locations.
- There are many benefits of planting early MG soybeans including, but not limited to, earlier harvest timing, earlier cover crop seeding, and risk management benefits.
- The objective of this trial was to determine the yield impact of early MG soybean product selection against the normal MG products for the location.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Fonda, IA	Silty clay loam	Corn	No tillage	5/25/18	10/19/18	65	140K
Storm Lake, IA	Silty clay loam	Corn	Conventional	5/25/18	10/20/18	65	140K
Marble Rock, IA – North	Loam	Corn	Strip tillage	5/22/18	10/18/18	55	140K
Marble Rock, IA - South	Loam	Corn	Strip tillage	5/22/18	10/18/18	55	140K
Huxley, IA	Clay loam	Corn	No tillage	5/19/18	10/17/18	60	140K
Atlantic, IA	Silty clay loam	Corn	Conventional	5/29/18	10/16/18	65	140K
Shenandoah, IA	Silty clay loam	Corn	No tillage	5/31/18	10/15/18	60	140K
Victor, IA	Silty clay loam	Corn	Conventional	5/18/18	9/24/18	60	140K

- This trial was broken into two sets, North and South Iowa, with a total of eight locations – four locations in the north set and four locations in the south set:
 - North Set – Fonda, Storm Lake, Marble Rock North, and Marble Rock South
 - South Set – Huxley, Atlantic, Shenandoah, and Victor
- Each set consisted of 18 unique soybean products:
 - Nine products are considered early MG
 - North Set – 1.1 MG to 1.8 MG
 - South Set – 2.0 MG to 2.4 MG
 - Nine products are considered normal MG
 - North Set – 2.0 MG to 2.4 MG
 - South Set – 2.9 MG to 3.5 MG
 - The nine 2.0 to 2.4 MG products were the same products for both the north and south sets.
- The plots consisted of four, 15-ft-long rows in 30-in row spacing with three replications.
- The Shenandoah site exhibited above average levels of frog-eye leaf spot and insect feeding.
- Above average levels of sudden death syndrome were observed at the Victor site.
- The Marble Rock North site was impacted with hail on August 28.



Yield Observations When Shifting to Earlier Maturity Group Soybeans

Understanding the Results

- The effect of maturity group on soybean yield was variable and highly dependent on the location. For example, Victor saw an 8 bu/acre yield advantage with early MG products, whereas Huxley realized a 7 bu/acre advantage with normal MG products.
- In general, Atlantic, Victor and Storm Lake saw some level of yield advantage with early MG soybean products versus the other locations where normal MG products gained some yield advantage. However, average site performance across all locations was nearly similar at 58 bu/acre.

What does this mean for your farm?

- In general, early MG soybean products yield close to late MG products, especially when conditions are favorable.
- This trial experienced unfavorable growing conditions in the locations where the normal MGs succeeded, including:
 - Excessive rain, wind, and hail in season followed by weather-delayed harvest
 - Lower management (no R3 growth stage fungicide/insecticide application)




Figure 1. Average yields of the nine products in each MG range at each location. There was severe insect and frogeye leaf spot damage at Shenandoah and hail damage at Marble Rock North.

- Finding the proper genetic package for a maturity group is still critical when considering planting early soybeans.
- More research is needed in the genetic pipeline to better understand which soybean products are better suited for the south.
- It should be noted that a MG shift may not be right for every operation and that its benefits could be defined in terms other than yield.

Legal Statements

The information discussed in this report is from a multiple site, replicated demonstration. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. Bayer is a registered trademark of Bayer Group. © 2018 Bayer Group, All Rights Reserved. 181213100712 121718JMG





Management Practices for Optimizing Yield and Productivity in Corn

Trial Objective

- Optimizing farm productivity requires the efficient and sustainable use of farm inputs and management practices, such as the choice of corn product, seeding rate, soil fertility, seed treatment, and pest management, that ultimately determine profitability.
- This study was conducted to evaluate the economic impact of different management inputs and practices on corn yield and profitability.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Huxley, IA	Clay loam	Soybean	Strip tillage	04/30/2018	09/28/2018	225	33K, 38K

- Two corn products were used for this trial:
 - A 110-day relative maturity SmartStax® RIB Complete® corn blend product
 - A 114-day relative maturity SmartStax® RIB Complete® corn blend product
- Each product was planted at a regional standard rate of 33,000 (33K) seeds/acre and a higher rate of 38,000 (38K) seeds/acre.
- The trial was carried out in 30-inch row spacing, six rows per treatment, with two replications.
- Management practices that were tested: choice of corn product, seeding rate, seed treatment, additional nitrogen, and fungicide application. These practices were compared in incremental stair-step treatments (Table 1).

Table 1. Treatments used in the trial with their associated costs.

Treatments	Input	110 RM Cost (\$/acre)	114 RM Cost (\$/acre)
33K	33,000 seeds/acre	\$ -	\$ -
33K+Q	QuickRoots® Dry Planter Box Corn (Q), \$6.19/acre	\$6.19	\$6.19
33K+Q+N	Side dress 32% UAN at V5 growth stage (N)	\$16.54	\$16.54
33K+Q+N+F	Fungicide application at VT/R1 growth stage (F)	\$48.54	\$48.54
38K	Additional 5,000 seeds/acre	\$25.50	\$26.13
38K+Q	QuickRoots® Dry Planter Box Corn (Q), \$7.19/acre	\$32.63	\$33.26
38K+Q+N	Side dress 32% UAN at V5 growth stage (N)	\$42.98	\$43.61
38K+Q+N+F	Fungicide application at VT/R1 growth stage (F)	\$74.98	\$75.61



Management Practices for Optimizing Yield and Productivity in Corn

- All treatments received a maximum return to nitrogen (MRTN) rate of 140 lb of nitrogen/acre in the form of 32% urea ammonium nitrate (UAN) in the spring during the strip-till operation. In the “N” treatments, an additional 45 lb/acre of UAN was side dressed at the V5 growth stage.
- All seed was treated with the Acceleron® Seed Applied Solutions ELITE offering, consisting of fungicide, insecticide, and nematicide with Enhanced Disease Control (EDC) for the control of early- to mid-season diseases caused by *Fusarium*, *Rhizoctonia*, and *Colletotrichum*.
- In the “Q” treatments, QuickRoots® Dry Planter Box Corn, a microbial seed inoculant, was added as a dry planter box formulation for enhanced nutrient availability.
- In the “F” treatments, Delaro® 325 SC fungicide was applied at the VT/R1 growth stage.
- Minimal levels of gray leaf spot and northern corn leaf blight were observed at the trial site.

Understanding the Results

- The early RM product produced higher yields than the late RM product in all treatments except for the 38KQNF treatment (Figures 1 and 2).
- There was a minimal yield response to QuickRoots® Dry Planter Box Corn in the early RM product, but a 4-6 bu/acre yield improvement in the late RM product at both seeding rates (Figures 1 and 2).
- In general, for the early RM product, the addition of inputs did not substantially improve yields at the higher seeding rate. At the standard seeding rate, additional nitrogen produced the highest yield response (Figure 1).
- For the late RM product, yield increased incrementally with the addition of inputs at the higher seeding rate. At the standard seeding rate, QuickRoots® Dry Planter Box Corn and additional nitrogen improved yields but not fungicide (Figure 2).




Figure 1. Yield response of two corn products to production inputs at the standard seeding rate.



Management Practices for Optimizing Yield and Productivity in Corn




Figure 2. Yield response of two corn products to production inputs at the higher seeding rate.




Figure 3. Average yield and economic impact of corn products in response to production inputs. Corn price was set at \$3.25/bu.



Management Practices for Optimizing Yield and Productivity in Corn

- In both corn products, the higher seeding rate out-yielded the standard rate across all treatments (Figure 1 and 2).
- In terms of economic value, the early RM product was more profitable than the late RM product at all treatment levels, except when fungicide was applied at the higher seeding rate (38KQNF) (Figure 3).
- The most profitable treatment in this trial was the early RM product planted at the standard rate with QuickRoots® Dry Planter Box Corn and additional nitrogen (33KQN) (Figure 3).
- At each treatment level, the yield difference between the two seeding rates was substantial enough for the higher rate to be more profitable than the standard rate. This is true for both corn products (Figure 3).
- For the late RM product, the treatment with the most inputs (38KQNF) produced the highest yield and the highest profit (Figure 3).
- Profit per acre was calculated by multiplying total yield of the treatment by \$3.50 minus the inputs selected for each treatment.

What Does This Mean for Your Farm?

- Corn products respond differently to farm inputs. Environmental factors during the growing season highly affect the yield response to inputs.
- Inputs like nitrogen will continue to provide positive yield responses and economic gains if it is used within the MRTN range for the region.
- Yield response to fungicides can be highly variable and depends on the growing season. It's unclear why there wasn't a consistent yield response to fungicide in this trial as there were minimal levels of gray leaf spot and northern corn leaf blight at the trial site.
- Where feasible, growers are encouraged to plant more than one corn product. This provides a good risk management strategy for their operation. They should also have a discussion with their trusted agronomists on how well a corn product of interest performs under different growing conditions and management practices.

Legal Statements

The information discussed in this report is from a single-site, replicated demonstration trial. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship.

B.t. products may not yet be registered in all states. Check with your seed brand representative for the registration status in your state.

IMPORTANT IRM INFORMATION: RIB Complete® corn blend products do not require the planting of a structured refuge except in the Cotton-Growing Area where corn earworm is a significant pest. SmartStax® RIB Complete® corn blend is not allowed to be sold for planting in the Cotton-Growing Area. **See the IRM/Grower Guide for additional information. Always read and follow IRM requirements.**

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Ready technology contains genes that confer tolerance to glyphosate, an active ingredient in **Roundup® brand agricultural herbicides.** Agricultural herbicides containing glyphosate will kill crops that are not tolerant to glyphosate. Acceleron®, Asgrow and the A Design®, Asgrow®, DEKALB and Design®, DEKALB®, Delaro®, QuickRoots®, RIB Complete®, Roundup Ready®, Roundup® and SmartStax® are registered trademarks of Bayer Group. Herculex® is a registered trademark of Dow AgroSciences LLC. LibertyLink® and the Water Droplet Design® are trademarks of BASF Corporation. All other trademarks are the property of their respective owners. ©2018 Bayer Group, All Rights Reserved. 181022185634 102518TAM





Yield Response of Corn Products to Seeding Rate

Trial Objective

- Research has indicated that corn yield has a positive correlation with seeding rate until a threshold is reached, beyond which yield decreases. Defining the seeding rate threshold for each corn product is difficult as it's highly affected by management practices and the environmental conditions during the growing season.
- However, knowing the threshold is very critical, as it forms the basis upon which other management practices, such as nitrogen rate, are based.
- The objective of this study was to determine the yield response of corn products to different seeding rates.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Huxley, IA	Clay loam	Soybean	Strip tillage	04/30/2018	10/04/2018	240	32K, 36K, 40K

- Each corn product was planted at 32,000, 36,000, and 40,000 seeds/acre.
- The trial was carried out in 30-inch row spacing, 3 rows/seeding rate, and 200-ft long strips.
- An application of 140 lb/acre of 32% UAN was applied in the spring as informed by the Climate FieldView™ platform.
- Weed management consisted of a pre- and post-emergence program and was applied across all plots.

Understanding the Results

Table 1. Average performance of DEKALB® corn brand blends across all three seeding rates.

Corn Brand Blend	Harvest Population (seeds/acre)				Average Grain Moisture (%)	Average Yield (bu/acre)	Average Yield Ranking
	32K	36K	40K	Average			
DKC50-08RIB	29	33	35	32.3	17.1	213.7	17
DKC51-38RIB	28	33	38	33.0	16.7	217.2	15
DKC54-38RIB	32	33	37	34.0	17.2	233.2	6
DKC54-74RIB	31	35	40	35.3	16.1	224.9	12
DKC55-53RIB	31	36	39	35.3	16.7	212.7	18
DKC55-84RIB	33	36	40	36.3	16.2	213.7	16
DKC57-97RIB	28	35	38	33.7	16.9	228.4	11
DKC58-06RIB	32	37	39	36.0	17.4	231.7	8
DKC58-34RIB	31	34	40	35.0	17.9	233.3	5
DKC60-87RIB	32	35	38	35.0	18.3	236.3	4
DKC61-98RIB	31	36	37	34.7	17.2	229.6	10
DKC62-20RIB	31	36	38	35.0	17.1	224.1	13
DKC62-52RIB	31	35	37	34.3	17.5	237.4	3
DKC62-78RIB	32	37	39	36.0	17.0	222.6	14
DKC63-21RIB	31	33	40	34.7	17.0	229.7	9
DKC63-57RIB	28	36	36	33.3	18.8	239.9	2
DKC64-34RIB	33	33	38	34.7	19.8	233.0	7
DKC66-75RIB	28	31	35	31.3	19.7	247.6	1



Yield Response of Corn Products to Seeding Rate




Figure 1. Yield response of DEKALB® corn brand blends (100 RM – 108 RM) to seeding rate.



Figure 2. Representative ears of DEKALB® corn brand blends (100 RM–108 RM) at the different seeding rates.



Yield Response of Corn Products to Seeding Rate




Figure 3. Yield response of DEKALB® corn brand blends (110 RM – 116 RM) to seeding rate.



Figure 4. Representative ears of DEKALB® corn brand blends (110 RM–116 RM) at the different seeding rates.



Yield Response of Corn Products to Seeding Rate

What Does This Mean for Your Farm?

- Traditionally, we consider a 5 bu/acre yield response in a 4,000 seeds/acre increment to be economical.
- Ten out of 19 corn products had an economical response from a 32,000 to 36,000 seeds/acre increase in seeding rate.
- Only three out of 19 corn products had an economical response from a 36,000 to 40,000 seeds/acre increase in seeding rate.
- This is the first year on average that the jump from 36,000 to 40,000 seeds/acre wasn't economical.
 - Several factors from this season, including nitrogen leaching after 17 inches of rain in June, may have caused the lack of yield response in the higher seeding rates.
- Contact your local DEKALB® representative when making product seeding rate decisions.

Legal Statements

Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship.


B.t. products may not yet be registered in all states. Check with your seed brand representative for the registration status in your state.

SmartStax® multi-event technology developed by Monsanto Company and Dow AgroSciences.

IMPORTANT IRM INFORMATION: RIB Complete® corn blend products do not require the planting of a structured refuge except in the Cotton-Growing Area where corn earworm is a significant pest. SmartStax® RIB Complete® corn blend is not allowed to be sold for planting in the Cotton-Growing Area. See the IRM/Grower Guide for additional information. Always read and follow IRM requirements.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Ready technology contains genes that confer tolerance to glyphosate, an active ingredient in Roundup® brand agricultural herbicides. Agricultural herbicides containing glyphosate will kill crops that are not tolerant to glyphosate. FieldView™ is a trademark of The Climate Corporation. DEKALB and Design®, DEKALB®, RIB Complete®, Roundup Ready®, Roundup®, SmartStax® and VT Double PRO® are registered trademarks of Bayer Group. Herculex® is a registered trademark of Dow AgroSciences LLC. LibertyLink® and the Water Droplet Design® is a trademark of BASF Corporation. Respect the Refuge and Corn Design® and Respect the Refuge® are registered trademarks of National Corn Growers Association. Climate FieldView™ services provide estimates or recommendations based on models. These do not guarantee results. Consult your agronomist, commodities broker and other service professionals before making financial, risk management, and farming decisions. More information at <http://www.climate.com/disclaimers>. ©2018 Bayer Group. All rights reserved. 181108072704 110918TAM





Characterization of Corn Products for Their Response to Nitrogen Fertilization

Trial Objective

- Nitrogen fertilization is an integral part of corn production and is partly responsible for some of the drastic yield increases in corn production. Nitrogen is also the focus of a lot of the research in soil fertility and crop nutrition, with emphasis on finding the right rate, timing, source, and placement.
- It is important to understand how corn products respond to nitrogen fertilization. Knowing a corn product's demand for nitrogen may help avoid applying nitrogen that is not needed, which offers both an environmental and input cost benefit. Using technologies such as the Nitrogen Advisor from Climate FieldView™ may help farmers monitor and maintain the right nitrogen status throughout the growing season.
- The objective of this study was to characterize corn products for their sensitivity to different nitrogen rates. Rates were selected to induce both nitrogen stress and excess nitrogen.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Huxley, IA	Clay loam	Soybean	Strip tillage	4/30/2018	10/04/2018	225	34K
Marble Rock, IA	Silty loam	Soybean	Strip tillage	5/17/2018	10/16/2018	220	34K
Victor, IA	Silty clay loam	Corn	Conventional	5/02/2018	9/27/2018	240	35K

- The nitrogen rates tested were:
 - Low – 30 lb/acre for the corn-soybean rotations or 50 lb/acre for the corn-corn rotation
 - Medium – 160 lb/acre (Huxley only)
 - High – 230 lb/acre
- All nitrogen applications were made before planting.
- The trial was carried out in 30-inch row spacing, with four rows per treatment, and 45-ft long plots with two replications.
- Weed management consisted of a pre- and post-emergence program.

Understanding the Results

- There was clear indication of nitrogen extremes as shown in Figure 1. Plants in the low-rate treatment showed chlorotic symptoms due to nitrogen stress, whereas those in the other treatments did not.
- Generally, grain yields increased with increasing nitrogen rate such that the low treatment had the lowest yields and the high treatment had the highest yields, except for five products at Huxley, in which the medium rate out-yielded the high rate. Also, at Marble Rock, the low rate out-yielded the high rate in DKC51-38RIB brand blend (Figure 2–4).
- Even though yields were highest at the Victor location, the biggest nitrogen response was observed at Huxley, with an average of a 124 bu/acre yield difference between the low and high rates, followed by Victor with a 49 bu/acre difference and then Marble Rock with a 46 bu/acre difference.
- The medium nitrogen rate at Huxley helped to identify products with very different nitrogen requirements, such as DKC57-97RIB and DKC62-20RIB brand blends (Figure 2).



Characterization of Corn Products for Their Response to Nitrogen Fertilization



Figure 1. An aerial image of the field layout of the nitrogen trial at the Huxley, Iowa site. Attention is drawn to the difference in foliage color due to the different nitrogen rates. The photo was taken around the V12 growth stage.




Figure 2. Performance of DEKALB® brand blends in response to different nitrogen rates at Huxley, Iowa. The horizontal lines indicate the average performance of all the products for that nitrogen rate. Low p-value < 0.05, LSD = 36 bu/acre; medium p-value < 0.05, LSD = 62 bu/acre; high p-value = < 0.05, LSD = 45.



Characterization of Corn Products for Their Response to Nitrogen Fertilization




Figure 3. Performance of DEKALB® brand blends in response to different nitrogen rates at Marble Rock, Iowa. The horizontal lines indicate the average performance of all the products for that nitrogen rate. Low p-value < 0.05, LSD = 20 bu/acre; high p-value < 0.05, LSD = 44 bu/acre.




Figure 4. Performance of DEKALB® brand blends in response to different nitrogen rates at Victor, Iowa. The horizontal lines indicate the average performance of all the products for that nitrogen rate. Low p-value < 0.05, LSD = 11 bu/acre; high p-value < 0.05, LSD = 11 bu/acre.



Characterization of Corn Products for Their Response to Nitrogen Fertilization

What Does This Mean for Your Farm?

- A corn plant's yield response to nitrogen is a complex phenomenon and is substantially impacted by the weather during the growing season, the soil type, and the inherent soil fertility.
- Wet conditions in May and June at the Huxley location may have led to substantial nitrogen loss, which can increase stress on the plants. This may explain, in part, the very low yields observed at the low nitrogen rate treatment.
- The Victor location, on the other hand, presented a high inherent soil fertility which produced an average of 233 bu/acre with only 50 lb/acre of nitrogen. Such a field needs to be sustainably managed to avoid nitrogen loss to the water system.
- At the current market trend of \$0.23/lb of nitrogen and \$3.75/bu for corn, a minimum of 12.3 bu/acre is required to pay for the difference between the low and high nitrogen rates. Thus, all but one product, DKC51-38RIB brand blend at Marble Rock, were profitable at all locations. A minimum of 4.3 bu/acre was required between the medium and high rates at the Huxley location.
- Corn products respond differently to farm inputs and they should be tested on a small scale before they are deployed for the whole farm.

Legal Statements

The information discussed in this report is from a single site, non-replicated demonstration. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship.

B.t. products may not yet be registered in all states. Check with your seed brand representative for the registration status in your state.


SmartStax® multi-event technology developed by Monsanto Company and Dow AgroSciences.

IMPORTANT IRM INFORMATION: RIB Complete® corn blend products do not require the planting of a structured refuge except in the Cotton-Growing Area where corn earworm is a significant pest. SmartStax® RIB Complete® corn blend is not allowed to be sold for planting in the Cotton-Growing Area. See the IRM/Grower Guide for additional information. Always read and follow IRM requirements.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Ready technology contains genes that confer tolerance to glyphosate, an active ingredient in Roundup® brand agricultural herbicides. Agricultural herbicides containing glyphosate will kill crops that are not tolerant to glyphosate. DEKALB and Design®, DEKALB®, RIB Complete®, Roundup Ready®, Roundup®, SmartStax® and VT Double PRO® are registered trademarks of Bayer Group. Herculex® is a registered trademark of Dow AgroSciences LLC. LibertyLink® and the Water Droplet Design® is a trademark of BASF Corporation. Respect the Refuge and Corn Design® and Respect the Refuge® are registered trademarks of National Corn Growers Association. ©2019 Bayer Group. All rights reserved. The information discussed in this report is from a single-site, replicated demonstration trial. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

FieldView™ is a trademark of The Climate Corporation. Climate FieldView™ services provide estimates or recommendations based on models. These do not guarantee results. Consult your agronomist, commodities broker and other service professionals before making financial, risk management, and farming decisions. More information at <http://www.climate.com/disclaimers>. ©2018 Bayer Group. All Rights Reserved. 181212072552 121218TAM





Effects of Nitrogen Rate on Corn Yield Potential

Trial Objective

- This trial was designed to help determine optimum nitrogen rates for new corn products and to demonstrate the interaction of crop inputs with specific corn products. The summary contains two years of results for some products and also results from some products grown only in 2018.

Research Site Details

Year	Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
2017	Victor, IA	Silty clay loam	Soybean	Conventional	4/21/17	10/5/17	220-260	35K
2018	Victor, IA	Silty clay loam	Soybean	Conventional	4/25/18	10/3/18	220-260	35K

- Nitrogen (N) was applied at rates of 110, 140, and 170 total lb N/acre as anhydrous ammonia (NH₃) in the spring 5 to 9 days before planting.
- Corn products were planted into each NH₃ zone consisting of six 30-inch rows, approximately 370 ft. long.

Understanding the Results

- The two-year average yield for seven products in a corn-soybean rotation was greatest with 140 lb total N/acre.




Figure 1. Average yield response of seven DEKALB® corn brand blends at three nitrogen rates over two years.



Effects of Nitrogen Rate on Corn Yield Potential




Figure 2. Two-year average yield response of three individual DEKALB® brand blends at three nitrogen rates.




Figure 3. Two-year average yield response of four individual DEKALB® brand blends at three nitrogen rates.



Effects of Nitrogen Rate on Corn Yield Potential




Figure 4. Yield response of three individual DEKALB® brand blends at three nitrogen rates in 2018.

What Does This Mean for Your Farm?

- The crop response to nitrogen is dependent on several variables, such as corn product, soil, and environment.
- Low nitrogen rates can limit yield potential, while excess rates are not beneficial for additional crop response and reduce the return on investment.
- The economic return to each additional unit of nitrogen will depend upon grain price and nitrogen price.

Legal Statements

The information discussed in this report is from a single site, non-replicated demonstration. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.


Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship.

B.i. products may not yet be registered in all states. Check with your seed brand representative for the registration status in your state. SmartStax® multi-event technology developed by Monsanto Company and Dow AgroSciences.

IMPORTANT IRM INFORMATION: RIB Complete® corn blend products do not require the planting of a structured refuge except in the Cotton-Growing Area where corn earworm is a significant pest. SmartStax® RIB Complete® corn blend is not allowed to be sold for planting in the Cotton-Growing Area. See the IRM/Grower Guide for additional information. Always read and follow IRM requirements.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Ready technology contains genes that confer tolerance to glyphosate, an active ingredient in Roundup® brand agricultural herbicides. Agricultural herbicides containing glyphosate will kill crops that are not tolerant to glyphosate. Asgrow and the A Design®, DEKALB and Design®, Roundup Ready®, Roundup® and SmartStax® are registered trademarks of Bayer Group. Herculex® is a registered trademark of Dow AgroSciences LLC. LibertyLink® and the Water Droplet Design® is a trademark of BASF Corporation. Respect the Refuge and Corn Design® and Respect the Refuge® are registered trademarks of National Corn Growers Association. ©2019 Bayer Group. All rights reserved. 181130131212 120318JMG





Effect of Fungicide on Yield and Plant Health

Trial Objective

- The application of a fungicide can protect corn plants from foliar diseases and increase overall plant health, which can lead to increased grain yield.
- Yield increases observed from the application of a fungicide greatly depend on corn product selection, as individual products respond differently to a fungicide application. While fungicide is often used as a high-yield management strategy, it can also be used to protect the yield of corn products with poor plant and stalk health ratings.
- The objective of this trial was to evaluate the impact that a fungicide application has on corn yield and late-season plant health.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Fungicide Date	Planting Rate
Atlantic, IA (southern set)	Silty Clay Loam	Soybean	Conventional	4/27/2018	10/31/2018	7/21/2018	35,000
Huxley, IA (both northern and southern sets)	Clay Loam	Soybean	Strip Till	5/9/2018	10/17/2018	7/17/2018	34,000
Marble Rock, IA (northern set)	Loam	Soybean	Strip Till	5/18/2018	10/24/2018	7/30/2018	36,000
Storm Lake, IA (northern set)	Silty Clay Loam	Soybean	Fall Vertical	5/8/2018	10/26/2018	7/24/2018	39,000
Victor, IA (southern set)	Silty Clay Loam	Soybean	Conventional	4/30/2018	10/27/2018	7/18/2018	35,000

- Ten DEKALB® corn products were divided into two sets based on relative maturity, with the northern set being located at Marble Rock, Storm Lake and Huxley, and the southern set being located at Atlantic, Victor and Huxley.
- Plots were planted as strip trials at four locations, with Huxley being arranged as a small-plot trial.
- The trial was replicated by location.
- Staygreen and disease ratings were collected during the growing season, and stalk strength and intactness were collected at harvest.
- Each site was sprayed with Delaro™ 325 SC fungicide (12 oz/acre) with a ground sprayer at brown silk.

Table 1. DEKALB® corn brand blends used in the trial with their associated ratings for stalk strength, staygreen, and harvest appearance. Ratings shown are general product ratings from the seed guide.

Corn Product	Stalk Strength	Staygreen	Harvest Appearance
DKC50-08RIB	3	3	4
DKC51-38RIB	3	2	2
DKC54-38RIB	2	3	3
DKC57-97RIB	2	2	2
DKC58-06RIB	4	2	2
DKC60-88RIB	3	3	3
DKC62-20RIB	3	4	4
DKC62-53RIB	3	4	5
DKC63-21RIB	3	3	3
DKC64-35RIB	1	1	1



Effect of Fungicide on Yield and Plant Health

Understanding the Results




Figure 1. Yield of DEKALB® corn brand blends with and without fungicide.

- Across all corn products, spraying a fungicide offered a 13 bu/acre advantage vs. the unsprayed treatment. For this study, a 6.8 bu/acre response was considered a profitable response (\$24/acre cost for fungicide application with \$3.50 corn).
- Fungicide use also increased plant health, as the average staygreen and intactness ratings improved from 5 to 3 and 6 to 2, respectively, for the sprayed products compared to the unsprayed products (data not shown).
- Fungicide application had a minimal effect on grain moisture, with a 0.6% difference in moisture between the sprayed and unsprayed treatments.




Figure 2. Pictures of DKC62-53RIB brand blend taken on 9/11 (left) and at harvest (10/30) at Atlantic, IA.



Effect of Fungicide on Yield and Plant Health

What Does This Mean for Your Farm?

- The 2018 growing season saw a range of moisture and temperature extremes occur across Iowa. Generally, the research sites saw a wet June, a dry July, and a very wet late summer/harvest season. This led to high levels of stalk and plant health issues due to excess moisture, disease, and lack of nitrogen.
- Such conditions may explain why a fungicide application was profitable across nearly all corn products tested in 2018. While fungicides do not cure plant diseases, a timely application can prevent foliar diseases from infecting the upper canopy.
- The results of this study suggest that a healthier upper canopy lead to increased photosynthetic activity later in the growing season, which resulted in increased yield in corn products sprayed with fungicide. While plant health was notably improved by fungicide use, we did not observe dramatic differences in stalk health between sprayed and unsprayed corn products.
- This trial will be repeated in 2019, with more focus placed on potential stalk health benefits derived from applying fungicide.

Legal Statements

The information discussed in this report is from a multiple site, replicated demonstration. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship.

B.t. products may not yet be registered in all states. Check with your seed brand representative for the registration status in your state.

SmartStax® multi-event technology developed by Monsanto Company and Dow AgroSciences.

IMPORTANT IRM INFORMATION: RIB Complete® corn blend products do not require the planting of a structured refuge except in the Cotton-Growing Area where corn earworm is a significant pest. SmartStax® RIB Complete® corn blend is not allowed to be sold for planting in the Cotton-Growing Area. See the IRM/Grower Guide for additional information. Always read and follow IRM requirements.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Ready technology contains genes that confer tolerance to glyphosate, an active ingredient in Roundup® brand agricultural herbicides. Agricultural herbicides containing glyphosate will kill crops that are not tolerant to glyphosate. DEKALB and Design®, Delaro®, Roundup Ready®, Roundup® and SmartStax® are registered trademarks of Bayer Group. Herculex® is a registered trademark of Dow AgroSciences LLC. LibertyLink® and the Water Droplet Design® is a trademark of BASF Corporation. Respect the Refuge and Corn Design® and Respect the Refuge® are registered trademarks of National Corn Growers Association. ©2019 Bayer Group. All rights reserved. 181212075254 121318JMG





2018 FFA Fantasy Farming Challenge

Trial Objective

- Since 2016, the Bayer Learning Center in Huxley, Iowa has engaged the Iowa FFA community in a farming contest dubbed the Fantasy Farming Challenge. The program allows students to make key decisions for a plot of corn to produce the highest yield and/or the highest profit.
- The students select several real-life crop production management decisions, each with an associated cost. The Learning Center plants each FFA chapters' plot using the selected management inputs. FFA chapters are invited to the Learning Center to see their plot and learn about Bayer Crop Science and the opportunities within agriculture.
- At the end of the 2018 season, each plot was harvested and 1st, 2nd, and 3rd place prizes were awarded for the highest yield and/or the highest profit.
- In 2018, 25 chapters participated in the program.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Huxley, IA	Clay loam	Soybean	Strip tillage	Various	09/28/2018	225	Various

- Each chapters' management decisions are shown in Table 1.
- Each plot was planted in 30-inch row spacing, 6 rows per chapter, and 375-ft long strips.
- Weed management was the same across all plots and consisted of a pre- and post-emergence program.

Understanding the Results

- Congratulations to our 2018 winners (Figure 1).
 - **Yield Winners**
 - **1st** Wilton (\$1,000)
 - **2nd** Diamond Trail (\$500)
 - **3rd** South Winneshiek (\$250)
 - **Net Profit Winners**
 - **1st** Wilton (\$1,000)
 - **2nd** Diamond Trail (\$500)
 - **3rd** S.E. Polk (\$250)
 - **3rd** Gilbert (\$250)
- We look forward to another exciting program in 2019.



2018 FFA Fantasy Farming Challenge

Table 1. Management decisions of the 25 FFA chapters involved in the 2018 Fantasy Farming Challenge.

FFA Chapter	Corn Brand Blend Products	Seeding Rate (seeds/acre)	Planting Date	Nitrogen (lb/acre)	Side Dress	Starter Fertilizer	Soil Insecticide	Follar Fungicide	Total Cost (\$/acre) *	Yield Rank	Net Profit Rank
Wilton	DKC64-34RIB	37,500	Early	200	No	Yes	No	No	\$ 712.83	1	1
Diamond Trail	DKC58-06RIB	34,000	Mid	240	Yes	No	No	No	\$ 700.47	2	2
South Winneshiek	DKC61-86RIB	33,500	Mid	250	No	Yes	No	Yes	\$ 750.69	3	8
Davis County	DKC64-34RIB	32,500	Early	185	Yes	Yes	No	Yes	\$ 724.10	4	5
Corydon-Wayne	DKC64-34RIB	38,000	Mid	230	Yes	Yes	Yes	Yes	\$ 808.15	5	15
Gilbert	DKC60-87RIB	36,000	Early	180	Yes	Yes	No	No	\$ 692.06	6	4
SE Polk	DKC60-87RIB	33,000	Mid	200	Yes	Yes	Yes	No	\$ 689.86	7	3
Rock Valley	DKC64-34RIB	34,000	Mid	165	No	Yes	Yes	No	\$ 719.87	8	7
Collins-Maxwell	DKC63-21RIB	35,500	Mid	170	Yes	Yes	No	No	\$ 712.01	9	6
Westwood	DKC63-21RIB	35,500	Mid	200	No	Yes	No	Yes	\$ 747.96	10	10
Newton	DKC63-21RIB	34,000	Mid	170	Yes	No	No	Yes	\$ 725.50	11	9
CAM	DKC54-38RIB	35,000	Early	240	Yes	Yes	Yes	Yes	\$ 743.51	12	14
Linn-Mar	DKC64-34RIB	35,200	Mid	181	No	No	No	Yes	\$ 728.13	13	13
Shenandoah-Davis-Rodgers	DKC60-87RIB	31,000	Early	160	Yes	No	Yes	No	\$ 698.50	14	11
Kingsley-Pierson	DKC58-06RIB	34,000	Mid	180	Yes	No	No	Yes	\$ 694.88	15	12
AC-GC	DKC64-34RIB	35,000	Mid	175	Yes	Yes	No	Yes	\$ 743.96	16	19
Albia	DKC64-34RIB	32,000	Late	165	No	Yes	No	No	\$ 743.80	17	20
Audubon	DKC60-87RIB	32500	Mid	155	Yes	Yes	Yes	Yes	\$ 729.05	18	16
Roland Story	DKC64-34RIB	32500	Mid	160	Yes	Yes	No	Yes	\$ 728.04	19	18
ADM	DKC64-34RIB	33000	Mid	155	Yes	No	No	Yes	\$ 716.39	20	17
Charles City	DKC54-38RIB	34000	Mid	200	Yes	Yes	No	Yes	\$ 713.47	21	21
Missouri Valley	DKC58-06RIB	35000	Mid	140	No	No	Yes	Yes	\$ 686.81	22	22
Ballard	DKC54-38RIB	33000	Late	145	Yes	Yes	Yes	No	\$ 697.97	23	23
SE Warren	DKC64-34RIB	38000	Mid	132	Yes	Yes	Yes	Yes	\$ 709.07	24	24
GMG	DKC54-38RIB	34000	Mid	90	Yes	No	Yes	No	\$ 645.63	25	25

Early, mid, and late planting dates were on 4/28/2018, 5/10/2018, and 5/24/2018, respectively.

*Chapters did not pay any cash amount to participate in the program. All costs associated with the program were paid by Bayer Crop Science.



2018 FFA Fantasy Farming Challenge




Figure 1. Yield and profit results of the 2018 FFA Fantasy Farming Challenge at the Bayer Learning Center at Huxley, IA.

What Does This Mean for Your Farm?

- Each growing season presents its own unforeseen challenges that make some well-intended decisions fall short. We hope students appreciate the challenges our farmers face each year.

Legal Statements

The information discussed in this report is from a single-site, replicated demonstration trial. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

Asgrow and the A Design®, Asgrow®, DEKALB and Design® and DEKALB® are registered trademarks of Bayer Group. ©2018 Bayer Group, All Rights Reserved. 181130124533 113018TAM





Comparison of Row Spacing by Management Practice

Trial Objective

- Generation of farm revenue requires the optimization of production inputs in a sustainable manner. Over the years, advances in agronomic research, including crop protection, germplasm, nutrition, and equipment technologies, have benefited farmers with more inputs than ever before.
- Deployment of these inputs should be carefully evaluated for each operation to determine their effects on yield, farm revenue, and the environment.
- With the current commodity prices, some farmers contemplate cutting operation costs by eliminating some inputs, while others consider certain inputs to be key to their success if used in an integrated system for the crop.
- The objective of this study was to compare low- and high-input corn management practices in two row-spacing systems.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Huxley, IA	Clay loam	Soybean	Conventional	5/10/2018	10/4/2018	225	33K, 38K

Two management treatments were tested:

1. Standard Management
 - 33,000 seeds/acre seeding rate
 - 140 lb/acre nitrogen pre-planting
 2. Premium Management
 - 38,000 seeds/acre seeding rate (\$25.50/acre for the additional 5,000 seeds/acre)
 - 140 lb/acre nitrogen pre-planting
 - 40 lb/acre nitrogen side-dressed at the V6 growth stage (\$9.20/acre)
 - Delaro™ 325 SC fungicide application at the VT/R1 growth stage (\$22/acre)
- The two treatments were tested in both 20-inch and 30-inch row spacing.
 - A 113 RM and 114 RM corn product were used for this trial.
 - The trial was carried out on 10-ft x 225-ft long plots with two replications.
 - 32% UAN was used as the nitrogen source.
 - The same pre- and post-emergence weed management program was used in both treatments.



Comparison of Row Spacing by Management Practice

Understanding the Results

- For the standard treatment, the plant population was higher in the 20-inch spacing than in the 30-inch spacing. For the premium treatment, the plant population was higher in the 30-inch spacing (Table 1).
- There were very minor differences in grain moisture content between the treatments in both row spacings (Table 1).
- The premium treatment substantially out yielded the standard treatment in both row spacings (Figure 1).
- The 20-inch spacing out performed the 30-inch spacing across all treatments (Figure 1).

Table 1. Average agronomic response of the standard and premium management treatments in 20-inch and 30-inch row spacing. Early stand count was taken at the V4 growth stage. Harvest population was taken a few days before harvesting.

Row Spacing (inches)	Management Treatment	Early Stand Count (1000 seeds/acre)	Harvest Population (1000 seeds/acre)	Grain Moisture (%)
20	Standard	33.13	34.75	19.13
	Premium	37.50	37	19.78
30	Standard	32.94	32.75	19.70
	Premium	38.00	39.75	19.45




Figure 1. Average yield response of standard and premium management treatments in 20-inch and 30-inch row spacing.



Comparison of Row Spacing by Management Practice

What Does This Mean for Your Farm?

- At most production sites, 20-inch row spacing has been shown to be a better row spacing than 30-inch for corn production. Where equipment is available, 20-inch row spacing is recommended. In this trial, a yield advantage of 6 to 8 bu/acre was realized.
- In most corn operations, foliar fungicides, additional nitrogen, and a higher seeding rate often result in some form of yield increases. The question most often is whether the yield increases would be adequate to offset the cost of inputs. In this trial, at the current grain price of \$3.74/bu, a minimum of 15 bu/acre was required to pay for the additional inputs of the premium treatment. Thus, the premium treatment was profitable in both row spacings, generating 6 to 8 bu/acre in net gain over the standard treatment (Figure 1).
- Crop yield response to farm inputs can be highly variable, often impacted by the cropping sequence, environmental conditions during the growing season, and the selected germplasm. It is advisable that they be used in an integrated manner to optimize their synergistic effects. In this trial, for example, an increased seeding rate would require additional nitrogen to meet the plant demand.

Legal Statements

The information discussed in this report is from a single-site, replicated demonstration trial. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Asgrow and the A Design®, Asgrow®, DEKALB and Design®, DEKALB® and Delaro® are registered trademarks of Bayer Group. ©2018 Bayer Group. All Rights Reserved. 181210083838 121018TAM





Comparison of Corn Row Spacing and Seeding Rate - Storm Lake, IA

Trial Objective

- Several years of research have indicated that corn yield is positively correlated with plant population until a threshold is reached, beyond which yield decreases. Defining the population threshold for each corn product is difficult as it is highly impacted by several factors including row spacing, management practices, and the environmental conditions during the growing season.
- Adjusting row spacing is one method to spread plant spacing to maximize agronomics and plant-to-plant competition.
- The objective of this trial was to compare corn product yield at 20-inch and 30-inch row spacing at three seeding rates.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Storm Lake, IA	Silty clay loam	Soybean	No tillage	05/08/2018	09/28/2018	250	33K, 38K, 43K

- Four corn products (100 RM, 105 RM, 110 RM and 114 RM) were each planted at 33,000 (33K), 38,000 (38K), and 43,000 (43K) seeds/acre at both 20-inch and 30-inch row spacing.
- The trial was carried out in 10-ft-wide by 100-ft-long plots with two replications.
- A total fertilizer application consisted of 167-57-93-14-1 (N-P-K-S-Zn), of which 150 lb of nitrogen in the form of 32% UAN was applied in the spring.
- Weed management consisted of an early post-emergence program.
- No fungicide or insecticide were applied.

Understanding the Results

- Except for the 100 RM product, the seeding rate of 33K seeds/acre produced the highest yields in 20-inch row spacing (Figure 1), and 38K seeds/acre produced the highest yields in 30-inch row spacing (Figure 2).
- In both 20-inch and 30-inch row spacing, average yield (across all seeding rates) increased as the relative maturity of products increased, with up to a 40 bu/acre difference between the 100 RM and the 114 RM products in 20-inch row spacing (Figure 1) and a 27 bu/acre difference in 30-inch row spacing (Figure 2).
- Across all products, 20-inch row spacing substantially out-yielded 30-inch row spacing at all seeding rates (Figure 3). When averaged across all corn products, 33K seeds/acre was the highest yielding seeding rate in 20-inch row spacing and 38K seeds/acre was the highest yielder in 30-inch row spacing (Figure 3).



Comparison of Corn Row Spacing and Seeding Rate - Storm Lake, IA




Figure 1. Effects of seeding rate on corn product performance in 20-inch row spacing.




Figure 2. Effects of seeding rate on corn product performance in 30-inch row spacing.



Comparison of Corn Row Spacing and Seeding Rate - Storm Lake, IA




Figure 3. Effects of seeding rate and row spacing on corn product performance, averaged across all four corn products.

What Does This Mean for Your Farm?

- At most production sites, 20 inch rows has been shown to be a better row spacing than 30 inch rows for corn production. Where equipment is available, this option should be considered. In this trial, an average yield advantage of 15 bu/acre was realized with 20-inch row spacing (Figure 3).
- By rule of thumb, we consider a 6.25 bu/acre response in a 5K seeds/acre increment to be economical. Thus, 38K seeds/acre was only economical in the 105 and 110 RM products in 30-inch row spacing. All other configurations were most economical at the 33K seeds/acre seeding rate.
- The research site experienced a tremendous amount of rainfall during the growing season. Such growing conditions affects nutrient status and does not favor high populations, especially in narrow row spacing. This may be part of the reason for the poor performance of the 43K seeds/acre seeding rate across the products. However, this doesn't represent every year or what we should expect for a response next season.

Legal Statements

The information discussed in this report is from a single site, replicated demonstration. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. Asgrow and the A Design®, Asgrow®, DEKALB and Design® and DEKALB® are registered trademarks of Bayer Group. All other trademarks are property of their respective owners. ©2018 Bayer Group, All Rights Reserved. 181213100546 121918JMG





The Effects of Row Spacing and Seeding Rates on Corn Yield Potential

Trial Objective

- This trial was designed to provide farmers in southern Iowa row width comparisons (20- and 30-inch row width systems) on later maturity corn products in Iowa and to help determine the yield response of higher seeding rates within each row width system.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
2017 Victor, IA	Silty clay loam	Soybean	Conventional	4/22/17	10/9/17	250	33K, 38K, 43K
2018 Victor, IA	Silty clay loam	Soybean	Conventional	4/25/18	10/4/18	250	33K, 38K, 43K

- Five DEKALB® corn brand blends of 110 to 114 relative maturity were planted in two adjacent blocks at two different row spacings and at three different seeding rates within each row spacing:
 - 6-row, 30-inch row spacing planted at 33,000 (33K), 38,000 (38K), and 43,000 (43K) seeds/acre
 - 12-row, 20-inch row spacing planted at 33K, 38K, and 43K seeds/acre
- A variable row spacing Case IH® 1215 Early Riser® planter unit was used for all plantings at general planting depth settings.
- Both blocks received 150 lb/acre of anhydrous ammonia in the spring. Cultural practices were identical.
- Individual plots were approximately 200 feet long.

Understanding the Results




Figure 1. Average yields by row spacing and seeding rate of five DEKALB® corn brand blends in 2017 and 2018.



The Effects of Row Spacing and Seeding Rates on Corn Yield Potential




Figure 2. Yields of DEKALB® corn brand blends at each row spacing and seeding rate in 2018.

- Across all corn products, spraying fungicide offered a 13 bu/ac advantage vs the unsprayed treatment. For this study, a 6.8 bu/ac response was considered a profitable response (\$24/ac cost for fungicide application with \$3.50 corn).
- Fungicide use also increased plant health, as the average staygreen/intactness rating for the unsprayed products improved from 5 to 3 and 6 to 2 (respectively) when both compared to the products sprayed with fungicide.
- Fungicide application had a minimal effect on grain moisture, with a 0.6% difference in moisture between the sprayed and unsprayed.

Legal Statements

The information discussed in this report is from a single site, two-year demonstration. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship.

B.t. products may not yet be registered in all states. Check with your seed brand representative for the registration status in your state.

SmartStax® multi-event technology developed by Monsanto Company and Dow AgroSciences.

IMPORTANT IRM INFORMATION: RIB Complete® corn blend products do not require the planting of a structured refuge except in the Cotton-Growing Area where corn earworm is a significant pest. SmartStax® RIB Complete® corn blend is not allowed to be sold for planting in the Cotton-Growing Area. See the IRM/Grower Guide for additional information. Always read and follow IRM requirements.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Ready technology contains genes that confer tolerance to glyphosate, an active ingredient in Roundup® brand agricultural herbicides. Agricultural herbicides containing glyphosate will kill crops that are not tolerant to glyphosate. DEKALB and Design®, RIB Complete®, Roundup Ready®, Roundup® and SmartStax® are registered trademarks of Bayer Group. Herculex® is a registered trademark of Dow AgroSciences LLC. LibertyLink® and LibertyLink® and the Water Droplet Design® are trademarks of BASF Corporation. ©2019 Bayer Group. All rights reserved. 181210075544 121218JMG





Tillage Systems in Corn and Soybean Production

Trial Objective

- Tillage systems and operations have evolved over the years to meet specific production and/or environmental objectives. Considerations, such as soil and water conservation, input costs, labor efficiency, timing of tillage, crop rotation, soil health, short- and long-term land usage, crop nutrient management, and weed and pest management, are some of the things that drive tillage decisions on the farm.
- With improvements in tillage implements and herbicide technologies, farmers have access to an array of tillage options, ranging from conventional tillage to minimum tillage to no-till. Many farms do not use a single tillage type for all operations. Instead, a different tillage type is often deployed to meet the productivity requirement of each piece of land. Once decided, the piece of land is managed with that tillage type for several years.
- As such, it becomes necessary to periodically evaluate the continued suitability of tillage systems.
- The objective of this trial was to evaluate corn and soybean productivity responses to conventional and strip tillage systems.

Research Site Details

Location	2018 Crop	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Huxley, IA	Soybean	Clay loam	Corn	Strip tillage and conventional	5/17/2018	10/19/2018	60	140K
Huxley, IA	Corn	Clay loam	Corn	Strip tillage and conventional	5/9/2018	9/27/2018	225	34K

- A 112 RM corn product and a 2.4 MG soybean product were used in the trial.
- The corn trial was on 60 x 500-ft long plots. The soybean trial was on 60 x 350-ft long plots. The trial was planted in 30-inch row spacing with two replications.
- Conventional tillage consisted of a chisel plow followed by a soil finisher. The chisel plow consisted of a two-gang disk unit followed by ripping shanks that went about 18 inches deep followed by a set of chisels to smooth out the soil surface and incorporate residue. The soil finisher unit was comprised of a disk gang, a cultivator, and tine harrow units.
- Strip tillage was carried out in conjunction with liquid nitrogen application. The strip-till bar unit consisted of a no-till coulters in the front, followed by a liquid nitrogen knife, followed by a Vulcan strip-till unit comprised of row cleaners, no-till coulters that penetrated 2-3 inches deep and 7 inches wide, and a rolling basket to break any large soil clumps and smooth the soil surface for planting.
- All tillage operations were carried out in the spring.
- All corn treatments received 140 lb/acre of nitrogen pre-planting, followed by a side dress of another 40 lb/acre at the VT growth stage. 32% urea ammonium nitrate (UAN) was used as the nitrogen source.
- Weed management consisted of pre- and post-emergence programs in both crops.
- Conventional tillage was used at the research site in previous years.



Tillage Systems in Corn and Soybean Production



Figure 1. The tillage systems used in the corn and soybean trials. Strip tillage is shown on the left for corn (top) and soybean (bottom). Conventional tillage is shown on the right for corn (top) and soybean (bottom).

Understanding the Results

Table 1. Effects of two tillage systems on the agronomic performance of corn and soybean. The early stand count was taken at the V4 growth stage. Harvest population was measured a few days before harvesting. Corn was planted at 34,000 seeds/acre and soybean at 140,000 seeds/acre.

Crop	Tillage	Early Stand Count (1000 seeds/acre)	Harvest Population (1000 seeds/acre)	Grain Moisture (%)
Corn	Conventional	34.3	34.6	17.04
	Strip	33.8	33.4	17.01
Soybean	Conventional	105.6	93.5	12.35
	Strip	103.9	98.8	12.25

- In both crops, tillage did not have a major impact on stand establishment and plant population (Table 1).
- Grain moisture content was not affected by tillage in either crop (Table 1).
- There was a substantial yield difference between tillage systems in both crops, with strip tillage out-yielding conventional tillage (Figure 2).



Tillage Systems in Corn and Soybean Production




Figure 2. Average yield advantage of strip tillage over conventional tillage in corn and soybean production systems.

What Does This Mean for Your Farm?

- Conservation tillage practices, including strip tillage, allow for better water infiltration through the soil profile instead of the wash-off/run-off associated with conventional tillage systems. This improved soil moisture profile, along with its associated soil nutrients, could explain the difference in yield observed. This is especially true considering that the weather conditions at the research site during the trial were wet and rainy in May and June followed by a dry July.
- The advantages of strip tillage, such as improved soil heath, structure, and increased organic matter, cannot be the reason for the yield advantages realized as it takes several years for these soil characteristics to develop.
- It should be noted that crop yield response to tillage could be widely variable and site-specific, as impacted by environmental factors, soil type and drainage, and the cropping sequence. Thus, it requires multiple years of research to truly determine the productivity of tillage systems.
- Most tillage operations start in the fall after harvest and then are left to weather/over-winter before being finished off in the spring for planting. Due to environmental conditions, all tillage operations in this trial were carried out in the spring and thus may not fully reflect the exact effects of tillage on cropping systems. Therefore, this trial will be repeated in the coming years to determine the best tillage system for the site.
- Regardless of the crop chosen, the right tillage type should be the one that provides the best economic returns while still ensuring better environmental stewardship.

Legal Statements

The information discussed in this report is from a single-site, replicated demonstration trial. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. Asgrow and the A Design[®], Asgrow[®], DEKALB and Design[®] and DEKALB[®] are registered trademarks of Bayer Group. ©2018 Bayer Group. All Rights Reserved. 181205123920 120618TAM





Cover Cropping and Tillage Systems in Soybean Production

Trial Objective

- In northern geographies, it is more difficult to get a cover crop established with the shorter growing season. Selecting an early maturing soybean product may allow time for better cover crop establishment, but could this practice negatively impact the yield potential of the farming operation?
- Eliminating a tillage pass through the field is another cropping system decision, but is there a yield penalty associated with no tillage?
- The objective of this study was to evaluate different cropping systems that integrate no-till, conventional tillage, cover crops, and product maturity selection.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
Storm Lake, IA	Silty clay loam	Corn	Conventional & no tillage	5/24/18	9/17/18	70	140K

- Four cropping systems were evaluated:
 - System 1 – Early soybean maturity (1.1 MG), early cover crop establishment, and no tillage.
 - System 2 – Normal soybean maturity (2.4 MG), late cover crop establishment, and no tillage.
 - System 3 – Early soybean maturity (1.1 MG), no cover crop, and no tillage.
 - System 4 – Normal soybean maturity (2.4 MG), no cover crop, and conventional tillage.
- Plots were 20-ft wide and 340-ft long strip trials with five replications.
- Soybeans were planted into the cereal rye cover crop in the cover crop systems.
- Cereal rye was terminated with an early post-emergence herbicide program.
- All treatments were treated with the same late post-emergence herbicide program.

Understanding the Results



Figure 1. Field picture showing the established cereal rye cover crop used for the trial. Dense vegetation on the right is the early established cover crop and the sparse vegetation on the left is the late established cover crop. Picture was taken just before soybean planting.



Cover Cropping and Tillage Systems in Soybean Production




Figure 2. Average yields of four soybean cropping systems. Systems 1 and 2 are the cover crop trials with the early MG product (System 1) and normal MG product (System 2). System 3 is the early MG product in no-till, and System 4 is the normal MG product in conventional tillage.

- Figure 1 indicates that the time of cover crop establishment made a tremendous difference in cover crop biomass at the time of soybean planting.
- With the two cover crop systems, the early MG soybean product with early cover crop establishment (System 1) out-yielded the normal MG product and late cover crop establishment (System 2) (Figure 2).
- In this study, no tillage with an early MG soybean product (System 3) out-yielded conventional tillage with a normal MG soybean product (System 4).
- In all trials, each soybean product performed similarly across the systems; however, the early MG product (Systems 1 and 3) outperformed the normal MG product (Systems 2 and 4).

What Does This Mean for Your Farm?

- Choosing the proper genetics is the most vital component of any cropping system. In this trial, the early MG soybean product provided over a 2.5 bu/acre advantage over the normal MG product in the cover crop systems (Systems 1 and 2). Thus, if chosen properly, early-maturing soybeans could be a better fit in the cover crop system with little to no yield penalty.
- In this trial, no-till did not show any yield drag when compared to conventional tillage, thus saving money with less trips across the field. In some situations, no-till may provide a yield advantage in some years.

Legal Statements

The information discussed in this report is from a single site, replicated demonstration. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. Asgrow and the A Design®, Asgrow®, DEKALB and Design® and DEKALB® are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. ©2018 Bayer Group, All Rights Reserved. 181214101607 121718JMG





Using 2018 Corn Rootworm Beetle Counts to Help Evaluate the Risk of an Infestation for 2019

Trial Objective

- The monitoring of corn rootworm (CRW) beetle numbers in current corn and soybean fields can be used to help assess the potential risk of a CRW infestation reaching economic damage levels in corn and soybean fields during the next growing season.
- Use of this information may help guide decisions regarding management strategies including corn and soybean product selection.
- The objective of this project was to measure adult CRW population levels in corn and soybean fields in 2018 to assist in risk evaluation for 2019.

Research Site Details

Location	Soil Type	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield (bu/acre)	Seeding Rate (seeds/acre)
1499 fields	Drained or well drained	See Figure 1	Conventional	---	---	110-250	28K-36K

- One to four Pherocon® AM non-baited trapping sites were established at 1499 field locations across the corn growing areas of IA, IL, IN, OH, MI, WI, MN, ND, SD, NE, KS, MO, and CO (Figure 1, Top).
- The trapping sites were installed in the interiors of corn and soybean fields that encompassed a variety of crop and management histories (Table 1).
- The Pherocon® AM traps were refreshed at 5- to 10-day intervals for 2-8 consecutive weeks through CRW adult emergence, mating, and egg laying phases (late July through late September). Following each sampling interval, the counts of adult northern and western CRW beetles were recorded and used to calculate the average number of CRW beetles/trap/day by field.
- At the end of the collective sampling period, the maximum capture value for each field was determined and the data were used in further analyses.

Understanding the Results

Table 1. Location of 2018 CRW beetle monitoring fields by crop (top) and characterization of 2018 sampled fields by present crop and previous crop with average maximum daily captures for western and northern CRW beetles (bottom).

2017 Crop	Previous Crop	Number of Sampled Fields	Average peak number of Beetles/Trap/Day		
			Northern Corn Rootworm	Western Corn Rootworm	Total
Corn	Corn	181	0.27	3.85	4.11
Corn	Rotated	154	0.28	0.46	0.74
Corn	Not Specified	842	0.05	1.26	1.30
Total Corn	All Rotations	1177	0.18	1.78	1.97
Soybean	Corn	322	0.02	0.40	0.42
Corn and Soybean	All Rotations	1499	0.16	1.59	1.75



Using 2018 Corn Rootworm Beetle Counts to Help Evaluate the Risk of an Infestation for 2019




Figure 1. Average number of beetles per trap per day summarized by crop and rotation. Data in this graph is the result of field trials conducted on 1499 field plots in 10 different states in 2018.

Categories for CRW beetle counts are based on action thresholds (beetles/trap/day) suggested by Extension entomologists at the Universities of Illinois and Iowa State and provide economic damage (ED) potential for the following season.^{1,2}

- Less than 2 beetles indicate a low risk of ED.
- Greater than 1 beetle suggests a low risk for ED but could indicate populations are increasing.
- Greater than 2 beetles indicate ED is likely if control measures are not used.
- Control measures include CRW *Bacillus thuringiensis* (*B.t.*)-protected corn products or soil-applied insecticides.
- Greater than 5 beetles indicate ED is very likely and populations are expected to be very high.



Using 2018 Corn Rootworm Beetle Counts to Help Evaluate the Risk of an Infestation for 2019

2018 CRW Beetle Survey Data.

- CRW populations were variable across the corn growing area. This suggests that environment and management are factors in determining CRW pressure levels.
- 19% of corn fields had counts exceeding the action threshold of 2 beetles/trap/day (Figure 1).
- 11% of the corn fields were approaching action threshold levels (Figure 1).
- Corn followed by corn had higher average maximum daily counts than 1st-year corn (4.7 vs. 0.74 beetles/trap/day (Table 1).
- 39% of continuous corn fields exceeded the action threshold (Figure 1).
- Counts from soybean fields in IL and eastern IA were low (0.42 beetles/trap/day) (Table 1).
- The threshold was exceeded in 5% of all soybean fields sampled (Figure 1).
- Counts of 0 were recorded in 14% and 38% of corn and soybean fields, respectively (Figure 1).

2018 Data Interpolation (Figure 2).

- Point data were interpolated to estimate populations and relative risk at the landscape level.
- To account for variations in sampling density and distribution, interpolations were based on average maximum values calculated within a systematic grid applied to the estimation area.
- On a broad scale, CRW populations, and consequently risk potential, is elevated in corn fields across eastern and southwest NE, northeast CO, west KS, southeast SD, as well as northwest, central, and east central IA.
- Corn rootworm populations continue to be relatively low in many parts of ND, MO, IL, and southern WI; however, localized hot spots can be found every year.
- Notable CRW beetle presence in soybean fields was isolated to small areas in north central IL and northeast IA.




Figure 2. 2018 CRW trap counts taken from 1499 corn and soybean fields across the Corn Belt.



Using 2018 Corn Rootworm Beetle Counts to Help Evaluate the Risk of an Infestation for 2019

Comparison of 2017 vs. 2018 CRW Beetle Data (Figure 3)

- Absolute comparisons between 2017 and 2018 populations should be made with low confidence due to large differences in sampling intensity and distribution. However, trends may still be reliably identified.
- Areas with large populations (i.e. “hot spots”) are consistent from year to year. Populations appear to have grown in some areas (e.g. IA) while are dissipating in others (e.g. portions of IL and southern WI).




Figure 3. Corn root worm trap count from 1177 corn fields in 2018 (see the 2017 research report for previous CRW counts).

What Does This Mean for Your Farm?

- CRW pose a threat to yield and profit, making it a pest that cannot be ignored. University research has demonstrated that even a moderate level of CRW feeding can cause yield losses averaging 15% with losses up to 45% or more being possible.³
- In the absence of site-specific data, local/regional surveys may provide insight at the landscape level and can be used to make informed decisions regarding management and product selection decisions.
- Beetle numbers and infestation geographies change. Continue to monitor present and historical data to gain information regarding CRW infestation potential. Use this information to help prepare for the 2019 season by selecting *B.t.*-protected corn products to protect your risk of CRW larvae damaging roots the following year.



Using 2018 Corn Rootworm Beetle Counts to Help Evaluate the Risk of an Infestation for 2019

SOURCES

- ¹ Western corn rootworm. *Diabrotica virgifera virgifera* LeConte. Extension & Outreach. Department of Crop Sciences. University of Illinois. http://extension.cropsciences.illinois.edu/fieldcrops/insects/western_corn_rootworm.
- ² Hodgson, E. and Gassmann, A. 2016. Guidelines for using sticky traps to assess corn rootworm activity. Integrated Crop Management. Iowa State University. <https://crops.extension.iastate.edu/cropnews/2016/06/guidelines-using-sticky-traps-assess-corn-rootworm-activity>.
- ³ Evaluating corn rootworm risk and economic impact. 2017. Agronomic Spotlight. Monsanto Company.

Websites verified 11/9/17. 171106192900

Legal Statements

For additional agronomic information, please contact your local brand representative. The information discussed in this report is from a multiple-site demonstration trial. This informational piece is designed to report the results of this demonstration and is not intended to infer any confirmed trends. Please use this information accordingly.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Asgrow and the A Design®, Asgrow®, DEKALB and Design® and DEKALB® are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. ©2018 Bayer Group, All Rights Reserved. 181127100429 010819HKG









Bayer Crop Science - Huxley Learning Center

1551 Hwy 210 | Huxley, IA 50124 | (515) 597-5900

www.bayer.com |  @TheHuxleyLC